



HERITAGE
Women's
CENTER

To Whom it May Concern:

Preventative dental cleanings and annual exams during pregnancy are recommended.

The second trimester is ideal for cavity fillings and crowns, which should be treated to reduce the chance of infection. Root canals or tooth extraction are sometimes emergent and necessary.

All elective procedures such as teeth whitening, cosmetic procedures and x-rays should be postponed until after the birth. When x-rays are necessary in emergent situations, use of a lead shield including an apron and thyroid collar is recommended.

If antibiotics are required, Penicillin, Amoxicillin, and Clindamycin are often used. Tetracycline and Fluoroquinolones should be avoided.

Anesthesia should be used at the smallest dose possible. Lidocaine is often used. It is recommended that elective procedures that require nitrous oxide use be avoided during pregnancy. However, some studies show that limited amounts of nitrous oxide may be acceptable when needed for extensive procedures. Narcotic pain medications are acceptable.

For any unanswered questions, please call our office at 629-255-2194 (222 Medical Office Building) or 629-255-2090 (Westgate Office).

Best,

Dr. Elizabeth Bleecker

Dr. Arielle Grand

Dr. Chibbi Iwelu

Dr. Nancy Lipsitz

Dr. Craig Martin

Dr. Bernadette Meadors

Dr. Lauren Pinkard

Dr. Erin Rebele

Dr. Anne Rossell

Dr. Niharika Saini

Dr. Allison Strnad